

Contents lists available at ScienceDirect

Combinatorial Chemistry - An Online Journal

journal homepage: www.elsevier.com/locate/comche

Combinatorial Chemistry Online Volume 16, Issue 4, April 2014

N.K. Terrett *


Ensemble Therapeutics Corp., Cambridge, MA 02139, USA

1. Current literature highlights


1.1. Solid-phase synthesis of peptide thioureas and thiazole-containing macrocycles

Thioureas occur in a number of biologically-active molecules, with a range of pharmacological properties including analgesic, anticancer, antifungal and antibacterial properties. In addition, thiourea derivatives are potent inhibitors of enzymatic targets for medical use. Thioureas can also be synthetically converted into thiazoles, a heterocycle that occurs both in nature and in many drug molecules. A recent paper has presented a novel method for the solid-phase synthesis of α -thiourea peptides, which were subsequently converted to thiazoles and incorporated into macrocyclic structures through ring-closing olefin metathesis [1].

The activation of *N,N'*-di-Boc-thiourea with CuSO_4 , HgCl_2 or Mukaiyama's reagent is a well-known method for the generation of guanidines. However, it was found that conditions could be modified such that the major product of the reaction of the thiourea with a solid-supported amino acid was the thiourea peptide. For example, phenylalanine immobilised on 4-hydroxymethylbenzoic acid (HMBA) resin (**1**) was treated with a pre-mixed solution in dry DMF of *N,N'*-di-Boc-thiourea (**2**), Mukaiyama's reagent (**3**) and triethylamine to give predominantly the thiourea derivative (**4**) rather than a guanidine derivative.




With conditions having been optimised for formation of α -thiourea peptides, the approach was successfully tested on the range of 20 proteinogenic amino acids, generating products in good yield irrespective of the size and nature of the side-chains. Deprotection of the thiourea (removing the Boc group) proceeded cleanly by treating with dilute stannic chloride in dichloromethane to give **5**, and thiazole formation was achieved by the Hantzsch reaction with α -haloketones to give **6**.



The scene was now set to attempt the formation of macrocycles containing thiazoles. A small collection of thiazoles was prepared using the protocol outlined above. In particular, cyclisation to the thiazole using bromopyruvic acid generated carboxylated thiazoles (**7**) which could undergo amide-forming reactions with amines following on-resin activation. Having already incorporated an allylglycine residue on the carboxy-terminal side of the peptide, coupling with an alkene on the carboxylated thiazole to give **8**, permitted macrocyclisation to product **9** by ring closing metathesis (RCM). The RCM reaction was found to proceed most successfully with the Hoveyda Grubbs second generation catalyst, generating a range of macrocycles with from 15 to 17 atoms. It was found that the reaction was sufficiently flexible to prepare any of four possible diastereoisomers, in excellent product yields and purities.

* Tel.: +1 617 492 6977; fax: +1 617 492 6689.

E-mail address: nterrett@ensembletx.com



Overall, this new methodology provides a mild and selective approach to *N*-terminally modified α -thiourea peptides on solid support. The approach has further been used to prepare a number of thiazoles and thiazole-containing macrocycles by the RCM reaction.

2. A summary of the papers in this month's issue

2.1. Polymer supported synthesis

No papers this month.

2.2. Solution-phase synthesis

Multi-component reactions (MCRs) feature the potential for abundant reagent combinations, inputs, and post-MCR modifications, which result in exceptionally high diversification power. An expeditious cascade protocol for the synthesis of functionalised imidazo[1,2-*a*]pyridin-3-ols has been developed based on the Patai reaction. With the availability of commercial reagents and high efficiency in expanding molecule diversity, this method offers advantages to the existing procedures for the synthesis of imidazo-pyridin-3-ol analogues [2].

A previously unknown class of highly substituted benzoquinoline-spirooxindoles has been easily prepared by a novel application of a mild and efficient catalyst, SbCl_3 , for carbon-carbon and carbon-nitrogen bond formation reactions. The approach reacted isatins and alkynes (dialkyl but-2-ynedioate) with an aromatic amine (2-naphthylamine) in a one-pot three-component reaction to give a library of dialkyl 2'-oxo-4*H*-spiro[benzo[*f*]quinoline-3,3'-indoline]-1,2-dicarboxylate derivatives in very good to excellent yields [3].

An efficient synthesis of diverse 2-aryl/heteroaryl substituted pyrimidinyl ethanones has been developed using a palladium-catalyzed Suzuki-Miyaura coupling reaction strategy. Use of $\text{Pd}(\text{OAc})_2$ in the presence of PPh_3 and Na_2CO_3 in 1,4-dioxane solvent was found to be the most effective reaction condition for this transformation [4].

The catalytic, highly regio-, diastereo-, and enantioselective synthesis of a small library of chiral spirooxindolopyrrolizidines has been described. This approach proceeds via a three-component 1,3-dipolar cycloaddition reaction of azomethine ylides, derived from isatin, with electron-deficient dipolarophiles, and 3-(2-alkenyl)-1,3-oxazolidin-2-ones. A chiral copper(II) complex of cyclohexane-1,2-bis(aryl)methyleneamine catalysed this process at room temperature [5].

A domino reaction for the rapid and diverse synthesis of spiro[1*H*-pyrazolo[3,4-*b*]benzo[*h*]dihydroquinolin-4,3-indolin-2-ones] has been reported. The synthesis represents a thermodynamically-favoured four-component reaction between phenylhydrazine, isatins, naphthylamines, and 3-ketoesters giving novel products in excellent yields under solvent-free conditions [6].

2.3. Scaffolds and synthons for combinatorial libraries

The efficient synthesis of novel glycoconjugate amino acid building blocks wherein the amino acid and carbohydrate moieties are linked via a sulphonamide functional group has been reported. The general reaction sequence consists of coupling a glycosyl thioacetate to an amino acid methyl ester followed by oxidation and deprotection of the carbohydrate moiety. The synthesis of derivatives from a range of amino acids has been demonstrated, with reaction at either the α -amino group of amino acid precursors or the sidechain ϵ -amino group of lysine precursors [7].

2.4. Solid-phase supported reagents

In a recent paper, a variety of ketoximes, prepared from the corresponding ketones, undergo Beckmann rearrangement using $\text{H}_3\text{PW}_{12}\text{O}_{40}$ (DTPA) in acetonitrile under reflux to yield *N*-substituted amides and lactams in good to excellent yields (72–96%). The method provides an efficient, clean, eco-friendly, and simple synthesis, using a catalyst which is cheap, moisture tolerant, recoverable, and reusable without much loss of its activity [8].

An efficient one-pot synthesis of 2-substituted quinoxalines from 1,2-diamines and phenacyl bromides has been developed using K10-montmorillonite (K10 clay) as a catalyst at 50 °C in acetonitrile. The method offers an easy route for the synthesis of substituted quinoxalines in high yields, and a plausible mechanism has been proposed. Furthermore, the K10 clay catalyst can be recovered by simple filtration and reused six times without any loss in its catalytic activity [9].

The development of sustainable catalysts that can combine advantages from both homogeneous and heterogeneous catalysis has raised considerable promise over the last decade. To this end, using ligation chemistries that show high functional tolerance, that are highly efficient, specific, versatile, and that require mild conditions are particularly favoured. Following these guidelines, the copper-catalysed azide-alkyne cycloaddition reaction (CuAAC) has been employed for the heterogenisation of homogeneous catalysts [10].

2.5. Novel resins, linkers and techniques

No papers this month.

2.6. Library applications

A four-component Ugi reaction (Ugi-4CR) using formylphenyl boronic acids under mild conditions has been developed for the synthesis of arylboronic acid analogues. The reactions were performed in methanol and accelerated by microwave irradiation, which makes this strategy suitable for constructing

boronic-containing chemical libraries. Two of the synthesised analogues were found to have cytotoxic activity against HepG2, MDA-MB231, and A549 cancer cell lines, demonstrating the potential application of this approach in developing novel boron-containing pharmaceuticals [11].

With a broad spectrum of antiviral activity against clinically relevant mutant viruses, analogues from a new pyridone series met a mandatory criterion for next generation NNRTIs. Using a simple synthetic approach, a library of inhibitors was constructed, leading to the identification of a new pyridone. In cell cultures, this new NNRTI shows a superior potency profile against a range of wild type and clinically relevant, resistant mutant HIV viruses [12].

In an effort to improve a therapy for giardiasis and amoebic dysentery, the creation of chemical libraries is necessary to obtain novel drugs with high activity combined with low toxicity. The synthesis of novel 3-tetrazolylmethyl-4H-chromen-4-ones via an Ugi-azide multicomponent reaction and their biological evaluation against *Entamoeba histolytica*, *Giardia lamblia* and *Trichomonas vaginalis* has been described. Biological results show that these compounds could be considered as candidates to anti-parasitic drugs, especially against *G. lamblia* [13].

Oroidin, (*E*)-*N*-(3-(2-amino-1*H*-imidazol-4-yl)allyl)-4,5-dibromo-1*H*-pyrrole-2-carboxamide, is a pyrrole alkaloid isolated from the marine sponge *Agelas oroides*. Routine screening in a panel of twelve cancer cell lines revealed the compound to be only poorly cytotoxic. The development of eight focused libraries comprising thirty compounds in total revealed two new compounds as potent inhibitors of cell growth in the panel of cell lines [14].

MurF ligase is a crucial enzyme that catalyses the ultimate intracellular step of bacterial peptidoglycan biosynthesis, and thus represents an attractive target for antibacterial drug discovery. The design, synthesis and evaluation of a new series of cyanothiophene-based inhibitors of MurF enzymes from *Streptococcus pneumoniae* and *Escherichia coli* have been described. Different structural modifications of the parent compounds resulted in a focused library of 37 new inhibitors, providing low micromolar inhibitors of MurF from *E. coli* and *S. pneumoniae* [15].

References

- [1] Cohrt AE, Nielsen TE. Solid-phase synthesis of peptide thioureas and thiazole-containing macrocycles through Ru-catalyzed ring-closing metathesis. *ACS Comb Sci* 2014;16(2):71–7.
- [2] Wang Y, Saha B, Li F, Frett B, Li H-y. An expeditious approach to access 2-arylimidazo[1,2-*a*]pyridin-3-ol from 2-amino pyridine through a novel Petasis based cascade reaction. *Tetrahedron Lett* 2014;55(7):1281–4.
- [3] Karmakar R, Kayal U, Bhattacharya B, Maiti G. One-pot three-component reaction for the synthesis of biologically important spiro[benzo[*f*]quinoline-3,3'-indoline] derivatives. *Tetrahedron Lett* 2014;55(7):1370–2.
- [4] Faldu VJ, Talpara PK, Shah VH. Efficient synthesis of diversely substituted pyrimidines by palladium catalyzed Suzuki-Miyaura coupling. *Tetrahedron Lett* 2014;55(8):1456–60.
- [5] Salahi F, Taghizadeh MJ, Arvinnezhad H, Moemeni M, Jadidi K, Notash B. An efficient, one-pot, three-component procedure for the synthesis of chiral spirooxindolopyrrolizidines via catalytic highly enantioselective 1,3-dipolar cycloaddition. *Tetrahedron Lett* 2014;55(9):1515–8.
- [6] Hosseinjani-Pirdehi H, Rad-Moghadam K, Youseftabar-Miri L. A four-component synthesis of novel spiro[pyrazoloquinoline-oxindoles] under solvent-free conditions. *Tetrahedron* 2014;70(9):1780–5.
- [7] Lopez M, Bornaghi LF, Poulsen S-A. Synthesis of sulfonamide-conjugated glycosyl-amino acid building blocks. *Carbohydr Res* 2014;386:78–85.
- [8] Kaur G, Rajput JK, Arora P, Devi N. Keggin-type Bronsted dodecatungstophosphoric acid: a quasi homogenous and reusable catalyst system for liquid phase Beckmann rearrangement. *Tetrahedron Lett* 2014;55(6):1136–40.
- [9] Jeganathan M, Dhakshinamoorthy A, Pitchumani K. One-pot synthesis of 2-substituted quinoxalines using K10-montmorillonite as heterogeneous catalyst. *Tetrahedron Lett* 2014;55(9):1616–20.
- [10] Fernandes AE, Jonas AM, Riant O. Application of CuAAC for the covalent immobilization of homogeneous catalysts. *Tetrahedron* 2014;70(9):1709–31.
- [11] Lian R-C, Lin M-H, Liao P-H, Fu J-J, Wu M-J, Wu Y-C, et al. Direct synthesis of the arylboronic acid analogues of phenylglycine via microwave-assisted four-component Ugi reaction. *Tetrahedron* 2014;70(9):1800–4.
- [12] Côté B, Burch JD, Asante-Appiah E, Bayly C, Bédard L, Blouin M, et al. Discovery of MK-1439, an orally bioavailable non-nucleoside reverse transcriptase inhibitor potent against a wide range of resistant mutant HIV viruses. *Bioorg Med Chem Lett* 2014;24(3):917–22.
- [13] Cano PA, Islas-Jácome A, González-Marrero J, Yépez-Mulia L, Calzada F, Gámez-Montaña R. Synthesis of 3-tetrazolylmethyl-4*H*-chromen-4-ones via Ugi-azide and biological evaluation against *Entamoeba histolytica*, *Giardia lamblia* and *Trichomonas vaginalis*. *Bioorg Med Chem* 2014;22(4):1370–6.
- [14] Dyson L, Wright AD, Young KA, Sako JA, McCluskey A. Synthesis and anticancer activity of focused compound libraries from the natural product lead, oroidin. *Bioorg Med Chem* 2014;22(5):1690–9.
- [15] Hrast M, Anderluh M, Knez D, Randall CP, Barreateau H, O'Neill AJ, et al. Design, synthesis and evaluation of second generation MurF inhibitors based on a cyanothiophene scaold. *Eur J Med Chem* 2014;73:83–96.

Further reading

Papers on combinatorial chemistry or solid-phase synthesis from other journals

- Yu G, Kuo D, Shoham M, Viswanathan R. Combinatorial synthesis and in vitro evaluation of a biaryl hydroxyketone library as antiviral agents against MRSA. *ACS Comb Sci* 2014;16(2):85–91.
- Dasgupta S, Kitov PI, Sadowska JM, Bundle DR. Discovery of inhibitors of shiga toxin type 2 by on-plate generation and screening of a focused compound library. *Angew Chem Int Ed* 2014;53(6):1510–5.
- Cai H, Sun Z-Y, Chen M-S, Zhao Y-F, Kunz H, Li Y-M. Synthetic multivalent glycopeptide-lipopeptide antitumor vaccines: impact of the cluster effect on the killing of tumor cells. *Angew Chem Int Ed* 2014;53(6):1699–703.
- Selvam R, Rohini KC, Arunan C, Subashchandran KP. Synthesis of biologically important angiotensin-II and angiotensin-IV peptides by using 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)phenoxy resin. *Asian J Chem* 2013;25(15):8673–6.
- Hansen MR, Le Quement ST, Jakobsen TH, Skovstrup S, Taboureau O, Tolker-Nielsen T, et al. Solid-phase synthesis and biological evaluation of N-dipeptido-homoserine lactones as quorum sensing activators. *ChemBioChem* 2014;15(3):460–5.
- Wu T, Pan L, Xu X, Liu Q. Regiodivergent heterocyclization: a strategy for the synthesis of substituted pyrroles and furans using α -formyl ketene dithioacetals as common precursors. *Chem Commun* 2014;50(15):1797–800 (Cambridge, England).
- Hurevich M, Seeberger PH. Automated glycopeptide assembly by combined solid-phase peptide and oligosaccharide synthesis. *Chem Commun* 2014;50(15):1851–3 (Cambridge, UK).
- Kandasamy J, Schuhmacher F, Hahn HS, Klein JC, Seeberger PH. Modular automated solid phase synthesis of dermatan sulfate oligosaccharides. *Chem Commun* 2014;50(15):1875–7 (Cambridge, UK).
- Thaner RV, Eryazici I, Farha OK, Mirkin CA, Nguyen SBT. Facile one-step solid-phase synthesis of multitopic organic-DNA hybrids via "click" chemistry. *Chem Sci* 2014;5(3):1091–6.
- Bak A, Daszykowski M, Kaminski Z, Kiec-Kononowicz K, Kuder K, Fraczyk J, et al. Probing an artificial polypeptide receptor library using a series of novel histamine H3 receptor ligands. *Comb Chem High Throughput Screen* 2014;17(2):141–56.
- Medina-Franco JL, Martinez-Mayorga K, Meurice N. Balancing novelty with confined chemical space in modern drug discovery. *Expert Opin Drug Discov* 2014;9(2):151–65.
- Wu J, Zhang Y, Maida LE, Santos RG, Welmaker GS, LaVoi TM, et al. Scaold ranking and positional scanning utilized in the discovery of nAChR-selective compounds suitable for optimization studies. *J Med Chem* 2013;56(24):10103–17.
- Sproviero M, Rankin KM, Witham AA, Manderville RA. Utility of 5'-O-2,7-dimethylpityl for solid-phase synthesis of oligonucleotides containing acid-sensitive 8-aryl-guanine adducts. *J Org Chem* 2014;79(2):692–9.
- El-Faham A, Khatib SN, Subiros-Funosas R, Albericio F. BOP-OMe, BOP-OBt, and BOP-OAc: novel organophosphonic coupling reagents useful for solution and solid-phase peptide synthesis. *J Pept Sci* 2014;20(1):1–6.
- Saiz C, Castillo V, Fontan P, Bonilla M, Salinas G, Rodríguez-Haralambides A, et al. Discovering *Echinococcus granulosus* thioredoxin glutathione reductase inhibitors through site-specific dynamic combinatorial chemistry. *Mol Divers* 2014;18(1):1–12.
- Shen Y, Wang J-Y, Song G-H. Ionic liquid-supported synthesis of piperazine derivatives as potential insecticides. *Mol Divers* 2014;18(1):195–202.
- Bennett CS. Principles of modern solid-phase oligosaccharide synthesis. *Org Biomol Chem* 2014.
- Hussain MK, Ansari MI, Kant R, Hajela K. Tandem C-2 functionalization-intramolecular azide-alkyne 1,3-dipolar cycloaddition reaction: a convenient route to highly diversified 9*h*-benzo[*b*]pyrrolo[1,2-*g*][1,2,3]triazolo[1,5-*d*][1,4]diazepines. *Org Lett* 2014;16(2):560–3.
- Collins JM, Porter KA, Singh SK, Vanier GS. High-efficiency solid phase peptide synthesis (HE-SPPS). *Org Lett* 2014;16(3):940–3.